

Consent for Bilateral Simultaneous Refractive Surgery

While many patients choose to have both eyes treated at the same surgical setting, there may be risks associated with simultaneous treatment that are not present when the eyes are treated on different days. **If you elect to have surgery performed on both eyes at the same time, you should understand both the possible advantages and disadvantages of your decision.**

Safety: The risks of infection, severe inflammation, delayed clouding of the cornea, corneal scarring and internal bleeding or retinal damage is very rare but potentially devastating. If these complications occur in one eye, they may also occur in the other. Should any of these complications happen, you could experience significant loss of vision or even temporary or permanent legal blindness. By choosing to have the procedure performed on separate days, you avoid the risk of having one or more of these complications in both eyes at the same time.

Accuracy: If there is an over-correction or under-correction in one eye, chances are it may happen in both eyes. If a retreatment is required in one eye, it is quite possible that your fellow eye may also require a retreatment. By having surgery on separate days, the doctor can monitor the healing process and visual recovery in the first eye and may be able to make appropriate modifications to the treatment plan for the second eye. In some patients, this might improve the accuracy of the result in the second eye. By correcting both eyes simultaneously, there is no opportunity to learn from the healing patterns of the first eye before treating the second eye.

Visual Recovery: Most patients experience rapid visual recovery, but some may experience symptoms such as blurred vision, night glare or ghost images that can result in prolonged recovery of normal vision. Blurred vision may rarely continue for several weeks, which could make driving difficult or dangerous and could interfere with your ability to work if it occurs in both eyes. There is no way of predicting how long your eyes will take to heal. If the eyes are operated separately, you can generally function with the fellow eye while the first eye fully recovers. However, there may be a period of imbalance in vision between your two eyes, producing a form of double vision. If you are able to wear a contact lens in your unoperated eye, the corrective lens could minimize this imbalance. The balance in vision between your two eyes will usually be restored more rapidly if they are operated on the same day.

The healing corneal flap is most susceptible to trauma during the first several weeks after surgery. Should both flaps become accidentally displaced, significant visual loss in both eyes may result.

Satisfaction: Both eyes tend to experience similar side effects. If you experience undesirable side effects such as glare, ghost images, increased light sensitivity, or corneal haze in one eye, you will likely experience them in both eyes. These side effects may cause a decrease in vision or other negative effects, and some patients have elected to not have their second eye treated. By having each eye treated on separate dates, you will have the opportunity to determine whether the procedure has produced satisfactory visual results without loss of vision or other uncommon undesirable side effects. If you are over age 40, you will also have an opportunity to experience the change in your close vision that results from the correction of your nearsightedness or farsightedness. This could influence your decision on whether or not to fully correct your other eye to maintain some degree of close vision without the need for glasses (monovision).

Convenience: It may be inconvenient for you to have each eye treated at separate visits because it would necessitate two periods of recovery from the laser surgery and might require additional time away from work.

Cost: Professional and facility fees may be greater if the eyes are operated on different days, and the additional time off work that may also be needed can be costly.

Consent Statement:

“I have read and understand the above risks and benefits of bilateral simultaneous refractive surgery, and I understand that this summary does not include every possible risk, benefit and complication that can result from bilateral simultaneous refractive surgery. My doctor has answered all of my questions about the procedure. I wish to have both of my eyes treated during the same treatment session if my doctor determines that the treatment in the first eye appeared to be technically satisfactory.

The reason(s) I wish to have both eyes treated at the same time are:

- Greater Convenience
- Possible faster recovery
- Less time away from work
- Contact lens intolerance and/or difficulty wearing contacts
- Elimination of possible vision imbalance between treated and untreated eyes
- Other _____

Patient Signature

Date

Printed Signature

Witness

Date

Reviewed risks, benefits and alternatives; all questions answered, patient understands and wishes to proceed.

Doctor Signature Date

IntraLASIK Correction Of Nearsightedness, Farsightedness, and Astigmatism Using IntraLase™ Technology

INDICATIONS AND PROCEDURE

This information is being provided to you so that you can make an informed decision about IntraLASIK, or “all-laser” LASIK. Traditional LASIK surgery involves two procedures: first, a microkeratome blade is used to create a flap on the cornea to expose the underlying tissue. After the flap is created, an excimer laser is used to reshape the eye by removing ultra-thin layers from the cornea in order to reduce farsightedness, nearsightedness, or astigmatism. Finally, the flap is returned to its original position, without sutures.

The IntraLASIK surgery also involves two procedures. First, instead of a microkeratome blade, it uses the FDA-approved IntraLase™ laser to create a flap with laser energy. The IntraLase laser is capable of creating extremely precise flaps by producing tiny bubbles inside the cornea that are 1/10,000 of an inch in diameter. The laser beam cannot penetrate into the eye beyond the cornea. After the flap is created, an excimer laser is used to reshape the eye by removing ultra-thin layers from the cornea in order to reduce farsightedness, nearsightedness, or astigmatism. The flap is returned to its original position, without sutures.

ALTERNATIVES

IntraLASIK is an elective procedure: there is no emergency condition or other reason that requires or demands that you have it performed. There are alternatives to this surgery: you could continue wearing contact lenses or glasses and have adequate visual acuity. There are also other types of refractive surgery, including LASIK with a microkeratome.

RISKS AND COMPLICATIONS

This procedure, like all surgery, presents some risks, many of which are listed below. You should also understand that there may be other risks not known to your doctor, which may become known later. Despite the best of care, complications and side effects may occur; should this happen in your case, the result might be affected even to the extent of making your vision worse. In giving my permission for IntraLASIK, I understand the following: The surgeon will use the FDA-approved IntraLase™ laser to create a flap, and then an FDA-approved excimer laser to reshape the eye. The long-term risks and effects of IntraLASIK are unknown. I have received no guarantee as to the success of my particular case. I understand that the following risks are associated with the procedure:

A. VISION-THREATENING COMPLICATIONS

1. I understand that the IntraLase™ laser or the excimer laser could malfunction, requiring the procedure to be stopped before completion. Depending on the type of malfunction, this may or may not be accompanied by visual loss.
2. I understand that irregular healing of the flap could result in a distorted cornea. This would mean that glasses or contact lenses may not correct my vision to the level possible before undergoing IntraLASIK. If this distortion in vision is severe, a partial or complete corneal transplant might be necessary to repair the cornea.
3. I understand that mild or severe infection is possible. Mild infection can usually be treated with antibiotics and usually does not lead to permanent visual loss. Severe infection, even if

successfully treated with antibiotics, could lead to permanent scarring and loss of vision that may require corrective laser surgery or, if very severe, corneal transplantation or even loss of the eye.

4. I understand that I could develop keratoconus. Keratoconus is a degenerative corneal disease affecting vision that occurs in approximately 1/2000 in the general population. While there are several tests that suggest which patients might be at risk, this condition can develop in patients who have normal preoperative topography (a map of the cornea obtained before surgery) and pachymetry (corneal thickness measurement) . Since keratoconus may occur on its own, there is no absolute test that will ensure a patient will not develop keratoconus following laser vision correction. Severe keratoconus may need to be treated with a corneal transplant while mild keratoconus can be corrected by glasses or contact lenses.
5. I understand that other very rare complications threatening vision include, but are not limited to, corneal swelling, corneal thinning (ectasia), appearance of “floaters” and retinal detachment, hemorrhage, venous and arterial blockage, cataract formation, total blindness, and even loss of my eye.

B. NON-VISION-THREATENING SIDE EFFECTS

1. I understand that there is a greater chance that the whites of my eyes may temporarily appear pink or red for several days to several weeks after surgery. This redness is more common with IntraLase-created flaps than with microkeratome-created flaps.
2. I understand that my vision after surgery using the IntraLase technology may not be clear immediately and that I might not notice improvement for several days to several weeks.
3. I understand that there may be increased sensitivity to light, glare, and fluctuations in the sharpness of vision. I understand these conditions usually occur during the normal stabilization period of from one to three months, but they may also be permanent.
4. I understand that there is an increased risk of eye irritation related to drying of the corneal surface following the LASIK procedure. These symptoms may be temporary or, on rare occasions, permanent, and may require frequent application of artificial tears and/or closure of the tear duct openings in the eyelid.
5. I understand that an overcorrection or undercorrection could occur, causing me to become farsighted or nearsighted or increase my astigmatism and that this could be either permanent or treatable. If permanent, I may need to use glasses or contact lenses. I understand an overcorrection or undercorrection is more likely in people over the age of 40 years and may require the use of glasses for reading or for distance vision some or all of the time.
6. After refractive surgery, a certain number of patients experience glare, a “starbursting” or halo effect around lights, or other low-light vision problems that may interfere with the ability to drive at night or see well in dim light. Although there are several possible causes for these difficulties, the risk may be increased in patients with large pupils or high degrees of correction. For most patients, this is a temporary condition that diminishes with time or is correctable by wearing glasses at night or taking eye drops. For some patients, however, these visual problems are permanent. I understand that my vision may not seem as sharp at night as during the day and that

I may need to wear glasses at night or take eye drops. I understand that it is not possible to predict whether I will experience these night vision or low light problems, and that I may permanently lose the ability to drive at night or function in dim light because of them. I understand that I should not drive unless my vision is adequate. These risks in relation to my particular pupil size and amount of correction have been discussed with me.

7. I understand that I may not get a full correction from my LASIK procedure and this may require future retreatment procedures, such as more laser treatment or the use of glasses or contact lenses.
8. I understand that there may be a “balance” problem between my two eyes after LASIK has been performed on one eye, but not the other. This phenomenon is called anisometropia. I understand this would cause eyestrain and make judging distance or depth perception more difficult.
9. I understand that, after LASIK, the eye may be more fragile to trauma from impact. Evidence has shown that, as with any scar, the corneal incision will not be as strong as the cornea originally was at that site. I understand that the treated eye, therefore, is somewhat more vulnerable to all varieties of injuries, at least for the first year following LASIK. I understand it would be advisable for me to wear protective eyewear when engaging in sports or other activities in which the possibility of a ball, projectile, elbow, fist, or other traumatizing object contacting the eye may be high.
10. I understand that there is a natural tendency of the eyelids to droop with age and that eye surgery may hasten this process.
11. I understand that there may be pain, irritation, or a foreign body sensation, particularly during the first 48 hours after surgery. I also understand that pain may be associated with complications such as infection.
12. I understand that temporary glasses either for distance or reading may be necessary while healing occurs and that more than one pair of glasses may be needed.
13. I understand that the long-term effects of LASIK are unknown and that unforeseen complications or side effects could possibly occur.
14. I understand that visual acuity I initially gain from LASIK could regress, and that my vision may go partially back to a level that may require additional surgery, or require glasses or contact lens use to see clearly.
15. I understand that the correction that I can expect to gain from LASIK may not be perfect. I understand that it is not realistic to expect that this procedure will result in perfect vision, at all times, under all circumstances, for the rest of my life. I understand I may need glasses to refine my vision for some purposes requiring fine detailed vision after some point in my life, and that this might occur soon after surgery or years later.
16. I understand that I may be given medication in conjunction with the procedure and that my eye may be patched afterward. I, therefore, understand that I must not drive the day of surgery and should not drive until I am certain that my vision is adequate for driving.



Refractive Surgery Financial Policy

The price that you pay for your refractive procedure is considered to be “all-inclusive” for up to one year after the date of your surgery. However, this does not include the cost of prescription eye drops. Additional charges will be incurred in the following situations:

Work-Up: If you (not your doctor) decide not to have the procedure after having the work-up, the charge for the work-up will be \$226. We can bill your insurance for the exam. If you cancel your surgery and decide to have the procedure done on a different date, any fees you incurred for the work-up will be deducted from the cost of your surgery.

- 1. Cancellation Fee:** If you opt to cancel within 48 hours of your surgery time, there will be a non-refundable charge of \$100.
- 2. Enhancements:** Enhancements are included in the cost of your procedure for up to one year if your visual acuity does not progress beyond 20/40, and you have attended all of your post-op appointments. If your visual acuity is better than 20/40 and you wish to have a retreatment, you will be charged for the enhancement.
- 3. Other medical conditions:** If during your post-operative period you develop other medical conditions (such as dry eye) you or your insurance company will be billed accordingly.

Though you are covered for one year post-operatively, your last scheduled “post-op” exam will be at 9-months. At one year post-op you will be due for a full eye exam, billable to insurance and subject to any co-pays that apply.

Monovision Trial:

The contact lens and fitting fees will be billed separately. There is no charge for trial lenses.

The fee for your procedure is \$___2500/eye_____ and is due on or before your procedure date, unless other arrangements have been made.

Other arrangements: \$200 off per eye if paid in full on surgery day.

Patient Signature _____ Date _____

Printed Name _____